



2017 HAWAII UNIVERSITY INTERNATIONAL CONFERENCES

SCIENCE, TECHNOLOGY & ENGINEERING, ARTS, MATHEMATICS & EDUCATION JUNE 8 - 10, 2017  
HAWAII PRINCE HOTEL WAIKIKI, HONOLULU, HAWAII

# AN ANALYSIS OF CAPITAL F IN THE FUNDAMENTAL THEOREM OF CALCULUS

GARRISON, JOSEPH M.

DEPARTMENT OF MATHEMATICS

MIDDLE GEORGIA UNIVERSITY

COCHRAN, GEORGIA.

Prof. Joseph M Garrison  
Department of Mathematics  
Middle Georgia State University  
Cochran, Georgia.

## **An Analysis of Capital F in the Fundamental Theorem of Calculus**

### **Synopsis:**

This paper will reflect on the integrating function central to the statement of the theorem which will be called Capital F. Capital F's relationship to all of the other antiderivatives of  $f(x)$  and the various structures of capital F producing the same function will be examined.

# An analysis of Capital F in the Fundamental Theorem of Calculus

## The Fundamental Theorem of Calculus

Suppose  $\int_c^x f(t)dt$  exists for all  $x, c \in [a, b]$ ;  $c$  is a constant real number.

Then if  $F_c(x) = \int_c^x f(t)dt$ , it follows that

$$\frac{d}{dx} F_c(x) = \frac{d}{dx} \int_c^x f(t) dt = f(x) \forall x, c \in [a, b]; c \text{ is a constant real number.}$$

After eluding Archimedes, Descartes, Barrow, Fermat and others ([https://en.wikipedia.org/wiki/History\\_of\\_calculus](https://en.wikipedia.org/wiki/History_of_calculus)) the fundamental theorem of calculus was finally discovered and validated through differentials and fluxions by Isaac Newton and later rigorously proved using limits by Augustin Louis Cauchy (1789--1857). "The modern proof of the Fundamental Theorem of Calculus was written in his Lessons Given at the École Royale Polytechnique on the Infinitesimal Calculus in 1823. Cauchy's proof finally rigorously and elegantly united the two major branches of calculus (differential and integral) into one structure."

<https://www3.nd.edu/~apilking/Math10550/Lectures/26.%20Fundamental%20Theorem.pdf>

This paper will reflect on the integrating function, which is central to the statement of the fundamental theorem of calculus. We will call this function Capital F. Capital F's relationship to all of the other antiderivatives of  $f(x)$  and the various structures of capital F producing the same function will be examined.

## Proposition One:

In other words, can all the antiderivatives of  $f(x)$  be generated by an appropriate choice of  $c$  in

$$\int_c^x f(t)dt \text{ where } x, c \in \mathfrak{R}?$$

This paper will show that in many cases it is not possible to generate all the antiderivatives of  $f(x)$  by an appropriate choice of  $c$ .

Note that we are not arguing that a given antiderivative does not exist, but that it cannot be generated by choosing a real number for the lower index

of the given  $F_c(x) = \int_c^x f(t)dt$  which we are calling Capital F.

## Proposition Two:

Suppose  $F_c(x) = \int_c^x f(t)dt$  and  $f(x)$  is continuous on  $\mathfrak{R}$ ;  $c$  is a constant.

Does there exist  $c_1 \neq c \in \mathfrak{R}$  such that

$$F_c(x) = \int_c^x f(t)dt = F_{c_1}(x) = \int_{c_1}^x f(t)dt?$$

In other words, are there different values of  $c$ , which produce the same F?

This paper will show that quite often there are many different values of  $c$  producing the same Capital F.

These values occur at the real zeros of Capital F!

Proposition One analyzed, first glance:

Can all the antiderivatives of  $f(x)$  be generated by an appropriate choice of  $c$  in

$$\int_c^x f(t)dt \text{ where } x, c \in \mathfrak{R} ?$$

$$\text{Let } F_c(x) = \int_c^x f(t)dt \text{ for a given } c \in \mathfrak{R}.$$

Now consider an antiderivative of  $f(x)$ , say  $G(x)$ ,  
obtained by choosing an appropriate  $k_1$  so that  $G(x) = F_c(x) + k_1$   
and  $G(x)$  has no real roots. This  $k_1$  exists when  $F_c(x)$  has either  
an upper bound or a lower bound.

Without loss of generality assume  $l$  is a lower bound.

$$\text{Choose } k_1 > |l|.$$

$$\text{Since } F_c(x) \geq l,$$

$$F_c(x) + k_1 > l + |l| > 0 \Rightarrow F_c(x) + k_1 > 0.$$

Then  $G(x)$  cannot be represented by

$$G_{c_2}(x) = \int_{c_2}^x f(t) dt, c_2 \in \mathfrak{R} \text{ because } G_{c_2}(c_2) = 0 \Rightarrow c_2 \text{ is a real root.}$$

But  $G(x)$  has no real root on  $\mathfrak{R}$ .

### Analysis Continued:

Let  $F_0(x) = \int_0^x t dt$  be a given antiderivative of  $f(x) = x$ .

$$F_0(x) = \frac{1}{2}x^2 \text{ because } F_0(x) = \int_0^x t dt = \frac{1}{2}t^2 \Big|_0^x = \frac{1}{2}x^2$$

(fundamental theorem of calculus).

Now any other antiderivative, say  $G(x)$ , can be

represented by  $G(x) = F_0(x) + k = \frac{1}{2}x^2 + k$  (mean value theorem).

Now what value of  $c$  in  $F_c(x) = \int_c^x t dt$

would produce the derivative  $\frac{1}{2}x^2 + k$ ?

$$\text{Since } \frac{1}{2}x^2 + k = F_c(x) = \int_c^x t dt = \frac{1}{2}x^2 - \frac{1}{2}c^2,$$

$$\text{then } -\frac{1}{2}c^2 = k \Rightarrow c^2 = -2k$$

$\Rightarrow k \leq 0$ . Therefore no antiderivative

of the form  $y = \frac{1}{2}x^2 + k; k \in \mathfrak{R}^+$

can be represented by  $\int_c^x t dt$ .

## Example for proposition One:

Suppose  $f(x) = x$ .

$G(x) = y = \frac{1}{2}x^2 + 4$  is a specific antiderivative for  $f(x)$ .

Let  $F(x) = \int_c^x t dt$

What real value for  $c$  above would generate  $G(x)$ ?

$F(x) = \int_c^x t dt$  is one antiderivative (fundamental theorem of calculus)

Any other antiderivative of  $x$  can be represented by

$\frac{1}{2}x^2 + k$  where  $k$  is a constant real number (mean value theorem).

$$F(x) = \int_c^x t dt = \frac{1}{2}t^2 \Big|_c^x = \frac{1}{2}x^2 - \frac{1}{2}c^2 = \frac{1}{2}x^2 + 4$$

The question becomes whether there exists  $c \in \mathfrak{R}$  such that

$-\frac{1}{2}c^2 = 4$  which would require that  $c^2 = -8$  and then  $c \notin \mathfrak{R}$ .

Any antiderivative of the form  $\frac{1}{2}x^2 + k$

must have  $k \leq 0$  to be represented by  $F(x) = \int_c^x t dt$ .

Therefore, any antiderivative of the form  $\frac{1}{2}x^2 + k$  where  $k$  is positive

cannot be represented by  $F(x) = \int_c^x t dt$ .

Let  $F_{c_i}(x) = \int_{c_i}^x f(t)dt$  be a given antiderivative of  $f(x)$  that is continuous on  $\mathfrak{R}$ .

$F_{c_i}(x) = \int_{c_i}^x f(t)dt$  exists by the fundamental theorem of calculus.

Now any other antiderivative, say  $G(x)$ , can be

represented by  $G(x) = F_{c_i}(x) + k = \int_{c_i}^x f(t)dt + k$  (mean value theorem).

$$F_{c_i}(x) = \int_{c_i}^x f(t)dt = F_{c_i}(x) - F_{c_i}(c_i).$$

Now what value of  $c$  in  $F_c(x) = \int_c^x f(t)dt$

would produce the antiderivative  $G(x) = F_{c_i}(x) + k$ ?

$$G(x) = F_{c_i}(x) + k = \int_c^x f(t)dt = F_{c_i}(x) - F_{c_i}(c) \Rightarrow$$
$$-F_{c_i}(c) = k$$

$$\text{and } F_{c_i}(c) + k = 0 = G(c)$$

$\Rightarrow$  that for the antiderivative  $G(x) = F_{c_i}(x) + k$

to be represented by  $\int_c^x f(t)dt$ , then the set containing the real roots

of  $G(x)$  must not be empty; and furthermore

all elements in this set produce

$$\text{identical } F_{c_i}(x) = \int_{c_i}^x f(t)dt .$$




## More Examples for Proposition One:

Antiderivatives with no real lower index in the generating function  $F(x) = \int_c^x t \, dt$  :

$$F_1(x) = \frac{1}{2}x^2 + 1$$

$$F_2(x) = \frac{1}{2}x^2 + 4$$

$$F_3(x) = \frac{1}{2}x^2 + 16$$



### Example:

Antiderivatives with real lower index in the generating function  $F(x) = \int_c^x t \, dt$  :


$$F_1(x) = \frac{1}{2}x^2 \quad \text{when } c=0$$

$$F_2(x) = \frac{1}{2}x^2 - 2 \quad \text{when } c = \pm 2$$

$$F_3(x) = \frac{1}{2}x^2 - 8 \quad \text{when } c = \pm 4$$

$$F_4(x) = \frac{1}{2}x^2 - 18 \quad \text{when } c = \pm 6$$

$$F_5(x) = \frac{1}{2}x^2 - 32 \quad \text{when } c = \pm 8$$



Proposition Two analyzed:

Suppose  $c_1 \neq c$  and  $\int_c^x f(t)dt = \int_{c_1}^x f(t)dt$ .

$$\int_c^x f(t)dt - \int_{c_1}^x f(t)dt = 0$$

$$\int_c^x f(t)dt + \int_x^{c_1} f(t)dt = 0$$

$$\int_c^{c_1} f(t)dt = 0 \Rightarrow$$

$c_1$  is a root of  $\int_c^x f(t)dt$ .

We also know that  $c$  is a root of  $\int_c^x f(t)dt$ ;

therefore the antiderivatives are = when the  $c_1$ 's are roots.

Suppose  $F_{c_1}(x) = \int_{c_1}^x f(t)dt$ .

Suppose  $c_2$  is a root of  $F_{c_1}(x)$ ,  $c_1 \neq c_2$

(note that  $c_1$  is also a root of  $F_{c_1}(x)$ ).

$$\text{Then } \int_{c_1}^{c_1} f(t)dt = 0$$

and  $F_{c_1}(c_2) = \int_{c_1}^{c_2} f(t)dt = 0$  because  $c_2$  is a root.

Consider  $x_1 \in$  the open interval  $(c_1, c_2)$ .

$$\int_{c_1}^{c_2} f(x) dx = \int_{c_1}^{x_1} f(x) dx + \int_{x_1}^{c_2} f(x) dx = 0$$

$$\text{or } \int_{c_1}^{x_1} f(x) dx = - \int_{x_1}^{c_2} f(x) dx = \int_{c_2}^{x_1} f(x) dx \Rightarrow$$

$$F_{c_1}(x_1) = F_{c_2}(x_1);$$

therefore  $F_{c_1}(x) = F_{c_2}(x)$ , and there are two distinct values of  $c$  producing the same Capital F.

## **Example for Proposition Two:**

Suppose  $F(x) = \int_1^x (2t - 3)dt$ .

Then  $F(1) = 0$ , and

all antiderivatives are of the form  $y = x^2 - 3x + c$ .

$$F(1) = 0 = 1 - 3 + c \Rightarrow c = 2,$$

so  $F(x) = x^2 - 3x + 2$ .

Note that  $f(2)=0$ .

$$F(x) = \int_1^x (2t - 3)dt = F(x) = \int_2^x (2t - 3)dt.$$

Thus, we have two values for the lower limit of integration producing the same  $F(x)$ .

These lower limits always occur at the real roots of

$$F(x) = x^2 - 3x + 2.$$

## The Connection between Proposition I and Proposition II:

Suppose there exists a real  $c$  such that

$F_c(x) = \int_c^x f(t)dt$ .  $F_c(x) = \int_c^x f(t)dt$  is an antiderivative of  $f(x)$ ,

continuous on  $\mathcal{R}$ (fundamental theorem of calculus).

Now any other antiderivative, say  $G(x)$ , can be represented by  $G(x) = F_c(x) + k$ (mean value theorem).

What value of  $c_1$  in  $F_{c_1}(x) = \int_{c_1}^x f(t) dt$

would produce the antiderivative  $G(x) = F_c(x) + k$ ?

$$G(x) = F_c(x) + k = \int_c^x f(t) dt + k = F_{c_1}(x) = \int_{c_1}^x f(t) dt = F_c(x) - F_c(c_1)$$

$$F_c(x) - F_c(c) + k = F_c(x) - F_c(c_1) \Rightarrow$$

$$F_c(x) - 0 + k = F_c(x) - F_c(c_1)$$

$$\text{and } F_c(c_1) = -k.$$

Thus for the antiderivative  $G(x) = F_c(x) + k$

to be represented by  $\int_{c_1}^x f(t) dt$ , the set containing the roots

of  $F_c(c_1) = -k$ , or  $G(x) = 0$  must not be empty; and furthermore,

all elements in this set are real numbers producing

identical functions  $F_c(x) = \int_c^x f(t) dt$ .

Given an antiderivative,  $F_c(x) = \int_c^x f(t)dt$ , of a continuous function and a desired constant of integration,  $k$ , in another antiderivative, in order to determine if the new antiderivative can be written in terms of Capital F with a real  $c$  solve the equation  $F_c(x) + k = \int_{c_1}^x f(t)dt = 0$

or solve  $G(x) = 0$ .

If there are real roots, it can be written in terms of Capital F with a real  $c$ .

The real roots all produce equal Capital F's.

**Example:**

Suppose  $F_c(x) = \int_c^x (2t + 1) dt$ .

$$F_0(x) = \int_0^x (2t + 1) dt =$$

$$(t^2 + t) \Big|_0^x = x^2 + x \text{ (fundamental theorem).}$$

Analyze  $x^2 + x - 6 = 0$


The roots  $\{-3, 2\}$  are values for  $c$  that

produce the same  $F_c(x) = \int_c^x (2t + 1) dt = x^2 + x - 6$ .

---

$$F_{2,-3}(x) = \int_2^x (2t + 1) dt = \int_{-3}^x (2t + 1) dt = x^2 + x - 6$$

---




Example:

Suppose  $F_c(x) = e^x + k$ .

What values of  $k$  allow  $F_c(x) = e^x + k$  to be written as  $\int_c^x e^t dt$ ?

Then  $e^x + k = 0$  when  $k < 0$ .

---



- Can be represented by real  $c$  in Capital F:  $y=e^x-2$
- Can be represented by real  $c$  in Capital F:  $y=e^x-3$
- Cannot be represented by real  $c$  in Capital F:  $y=e^x+2$



## Concluding Example

Note that any polynomial of odd degree can be represented by  $\int_c^x f(t)dt$ .

Suppose  $G(x) = x^3 - 2x^2 - x + 2$ .

What values of  $c$  allow  $G(x)$  to be written as  $F(x) = \int_c^x (3t^2 - 4t - 1)dt$ ?

Solve  $G(x) = x^3 - 2x^2 - x + 2 = 0$ .


The roots are  $\{-1, 1, 2\}$ .

$G(x) = x^3 - 2x^2 - x + 2 =$

$$\int_{-1}^x (3t^2 - 4t - 1)dt =$$

$$\int_1^x (3t^2 - 4t - 1)dt =$$

$$\int_2^x (3t^2 - 4t - 1)dt.$$



## Conclusion:

Let  $f(x)$  be the derivative of a differentiable function, say  $G(x)$ . The integrating functions

$F_{c_i}(x) = \int_{c_i}^x f(t)dt$  used in the Fundamental Theorem

are equal when the  $c_i$ 's are real roots of  $G(x)$ , and

$F_{c_i}(x) = \int_{c_i}^x f(t)dt = G(x)$ . There are many times when

an antiderivative of  $f(x)$  cannot be generated by

$c_i \in \mathfrak{R}$  in  $F(x) = \int_{c_i}^x f(t)dt$ . This occurs when  $G(x)$  has

no real root. The lower index  $c_i$  determines the

value of the constant of integration  $k$ , in the

antiderivative  $F(x) = G(x) + k$  where  $G(x)$  is any antiderivative because  $c_i$  is a root of  $F(x)$ . The final

concern is whether there is a representation of  $G(x)$

$= F_{c_i}(x) = \int_{c_i}^x f(t)dt$  where  $f(x)$  is not the derivative of

$G(x)$ . The fundamental theorem of calculus

guarantees that no such function exists.

QED